

A Current Assessment of the Nature of PM_{2.5} in Steubenville, Ohio, Using SCAMP Monitoring Data

Ohio Air Quality and Coal Research Symposium
Athens, Ohio
December 3, 2004

D. P. Connell, S. E. Winter, J. A. Withum

CONSOL Energy Inc. Research & Development
South Park, Pennsylvania



Why Steubenville?

- Most polluted of the Harvard Six Cities
 - Mean PM_{2.5} Concentration = 29.6 µg/m³, 1979-1985
 - Extensive PM_{2.5} data record
- Major changes have occurred
 - Steubenville-Weirton MSA lost 4,200 manufacturing jobs in 1990s (decline of steel industry)
 - Population decreased by 7.4% in 1990s
- Likely a nonattainment area under PM_{2.5} NAAQS



The Steubenville Comprehensive Air Monitoring Program (SCAMP)

- Two-year comprehensive program for monitoring PM_{2.5} and co-pollutants
- Steubenville, Ohio, and surrounding region
- May 2000 – May 2002
- Two major study components:
 - Indoor/Personal
 - Personal sampling of children and elderly volunteers
 - Indoor sampling in participants' homes
 - Outdoor
 - Participants' homes
 - Central site in Steubenville
 - Four remote sites located at cardinal compass points around Steubenville



SCAMP Outdoor Ambient Goals

- Compare urban PM_{2.5} concentration / composition with remote PM_{2.5} concentration / composition (determined using FRM)
- Study associations among PM_{2.5}, co-pollutants, and weather conditions
- Provide a comprehensive database for use in epidemiological and transport studies and in compliance program development



Steubenville Site



- PM_{2.5} FRM
 - Mass (1/1 days⁻¹)
 - Ions (1/4)
 - Elements in WS Fraction (1/4)
- PM_{2.5} Speciation Sampler
 - EC, OC (1/4)
 - Elements in Acid-Digestible Fraction (1/4)
- PM_{2.5} TEOM
 - Mass (continuous)
- PM₁₀ FRM
 - Mass (1/1)
 - Ions (1/4)
 - Elements in WS Fraction (1/4)




- FRM or FEM Gas Analyzers
 - SO₂, CO, NO_x, O₃ (continuous)
- 10-m Meteorological Tower
 - Weather Conditions (continuous)
- Burkard Volumetric Spore Trap
 - Pollen and Mold Spores (1/1)

Steubenville PM_{2.5} Concentration / Composition


May 2000 – May 2002



Seasonal Variability




Spatial Variability



Less Similar

Intersite Coefficients of Divergence Based Upon 21 PM_{2.5} Components




“Background” Sites

$$CD_{jk} = \sqrt{\frac{1}{p} \sum_{i=1}^p \left(\frac{x_{ij} - x_{ik}}{x_{ij} + x_{ik}} \right)^2}$$



Local Source Contributions

PM_{2.5} and Major Components



84 71 68 82



% ST > BG



Local Source Contributions

Elements in the Water-Soluble Fraction


	% ST > BG	Loc. (ng/m ³)	Loc. (% of BG)
Al	71	4.8	49
As	73	0.66	43
Ba	79	0.7	65
Cd	71	0.10	31
Ca	73	28	46
Co	43	-0.004	-10
Cu	59	0.7	31
Fe	71	11.8	106
Pb	69	3.1	78

	% ST > BG	Loc. (ng/m ³)	Loc. (% of BG)
Mg	79	18	145
Mn	83	4.5	154
Ni	47	0.1	11
K	59	16	21
Se	54	-0.44	-9
Na	73	19	32
Sn	38	-0.017	-9
V	63	0.44	66
Zn	75	25.5	140




Gas and Weather Percentiles On Highest / Lowest PM_{2.5} Days


Days with Highest 10% PM_{2.5} Concentrations




Days with Lowest 10% PM_{2.5} Concentrations



Seasonally Dependent Correlations



$R^2 = 0.03$
 $m = 0.16$




$R^2 = 0.52$
 $m = 0.53$




Diurnal Variability

PM_{2.5} (TEOM)





Gaseous Pollutants



PM_{2.5} Episode – Hourly Data


January 26 – January 30, 2002




PM_{2.5} Episode – Hourly Data

January 26 – January 30, 2002


Hourly Average PM_{2.5} and SO₂




Hourly Average PM_{2.5} and CO



Hourly Average PM_{2.5} and NO_x




Hourly Average PM_{2.5} and O₃




PM_{2.5} Episode – Hourly Data

July 30 – August 4, 2001


Hourly Average PM_{2.5} and Solar Radiation



Hourly Average PM_{2.5} and NO_x




Hourly Average PM_{2.5} and SO₂




PM_{2.5} Composition During Episodes

Ions, Carbon, Elements in Water-Soluble Fraction

Summertime Episode: 8/3/01 09:00 - 8/4/01 09:00



Wintertime Episode: 1/26/02 09:00 - 1/27/02 09:00



Summary

- Average $\text{PM}_{2.5}$ concentration in Steubenville has decreased by more than $10 \mu\text{g}/\text{m}^3$ since Six Cities Study; still more than $3 \mu\text{g}/\text{m}^3$ above annual NAAQS (based on 2000-2002 data)
- Sulfate (31.3 wt%) and organic material (25.0 wt%) are the major components of $\text{PM}_{2.5}$ in Steubenville
- Local sources on average contribute an estimated $4.6 \mu\text{g}/\text{m}^3$ to Steubenville's $\text{PM}_{2.5}$ concentration
 - Sulfate, Nitrate, and Ammonium account for about $2 \mu\text{g}/\text{m}^3$
 - Among elements within the water-soluble fraction, Mg, Mn, Zn, and Fe show the greatest percent urban increment
- $\text{PM}_{2.5}$ concentrations were positively associated with CO , NO_x , and SO_2 concentrations
 - Associations between $\text{PM}_{2.5}$ and some gaseous pollutants (i.e., NO_x and CO) were strongly dependent upon season



Summary

- PM_{2.5} exhibited a diurnal pattern similar to CO and NO_x; concentrations > 65 µg/m³ were never observed during the mid-afternoon
- High PM_{2.5} concentrations tended to occur on warm, high-pressure days; low concentrations tended to occur on cool, windy days with low solar radiation
- PM_{2.5} episode case studies:
 - Cool season (nocturnal temperature inversions)
 - PM_{2.5} concentrations showed strong diurnal variation
 - Strong associations among PM_{2.5}, NO_x, CO, SO₂
 - Enrichment: Zn, Mn, Mg, EC, Ca, Pb
 - Warm season
 - PM_{2.5} concentrations more chronically elevated
 - Associations among PM_{2.5} and gases not as strong
 - Enrichment: SO₄²⁻, NH₄⁺, Ni, Fe, Sn



Publications

- **Connell et al. (2004)** The Steubenville Comprehensive Air Monitoring Program (SCAMP): Overview and Statistical Considerations, *J. Air & Waste Manage. Assoc.*, in press.
- **Connell et al. (2004)** The Steubenville Comprehensive Air Monitoring Program (SCAMP): Associations Among PM_{2.5}, Co-Pollutants, and Meteorological Conditions, *J. Air & Waste Manage. Assoc.*, in press.
- **Connell et al. (2004)** The Steubenville Comprehensive Air Monitoring Program (SCAMP): Analysis of Short-Term and Episodic Variations in PM_{2.5} Concentrations Using Hourly Air Monitoring Data, *J. Air & Waste Manage. Assoc.*, in press.



Funding

U.S. DOE – National Energy Technology Laboratory

Ohio Coal Development Office

Electric Power Research Institute

American Petroleum Institute

National Mining Association

American Iron and Steel Institute

Edison Electric Institute

National Institute of Environmental Health Sciences

U.S. Environmental Protection Agency

CONSOL Energy Inc.



Participating Groups

CONSOL Energy Inc. R&D

Harvard School of Public Health

Ohio University

Franciscan University of Steubenville

Wheeling Jesuit University

St. Vincent College

Optimal Technologies

Air Quality Sciences, Inc.

Control Analytics, Inc.

